

Bose Condensate in External Potential: A Thomas-Fermi Approach¹

Kerson Huang

Center for Theoretical Physics, Laboratory for Nuclear Science
and Department of Physics
Massachusetts Institute of Technology
Cambridge, MA 02139

1 Thomas-Fermi Theory

The Thomas-Fermi method [1] [2] was designed for the calculation of the electron density in a heavy atom, by treating the electrons as locally free. Lieb and Simon [3] showed that the treatment is exact in the limit when the atomic number goes to infinity. Application to a confined Bose condensate was pioneered by Goldman, Silvera, and Legget [4], and by Oliva [5], and recently reconsidered by Chou, Yang, and Yu [6]. I shall describe some work on this subject, done in collaboration with E. Timmermans and P. Tommasini [7].

First, let us review the original method of Thomas and Fermi. Suppose $V(r) = -e\Phi(r)$ denotes the effective potential energy of an electron in an atom, at a distance r from the nucleus. (See Fig.1).




Figure 1: Potential energy of an electron in atom.

The condition that the electron is in a bound orbit is that

$$\frac{\hbar^2 k^2}{2m} + V(r) \leq 0 \quad (1)$$

¹Contribution to Workshop on Bose-Einstein Condensation, Institute for Theoretical Atomic and Molecular Physics, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, (August 19-30, 1996).

where k is the local wave vector. Assume that all available states are occupied. Then the local Fermi momentum $\hbar k_F(r)$ is given by

$$\frac{\hbar^2 k_F^2(r)}{2m} + V(r) = 0 \quad (2)$$

which gives

$$\hbar k_F = \sqrt{2me\Phi(r)} \quad (3)$$

We know that $k_F(r)$ is related to the local density $n(r)$ by

$$2(4\pi/3)k_F^3 = n(r) \quad (4)$$

where the factor 2 comes from spin. This relates the density to the potential $\Phi(r)$. We now use the Poisson equation

$$\nabla^2 \Phi(r) = -4\pi[Ze\delta(\mathbf{r}) - en(r)] \quad (5)$$

For $r \neq 0$ this equation is of the form

$$\nabla^2 \Phi(r) + C\Phi^{3/2}(r) = 0 \quad (6)$$

where C is a constant. One can solve this equation, and then obtain $n(r)$. A comparison between Thomas-Fermi and Hartree results for Rb is sketched in Fig.2.




Figure 2: Electron density in Rb: Comparison between Thomas-Fermi and Hartree approximations.

The essence of the method is that one assumes there is a local chemical potential $\mu_{\text{eff}}(r)$, related to the true chemical potential μ by

$$\mu_{\text{eff}}(r) = \mu - V(r) \quad (7)$$

In the earlier discussion, the chemical potential (Fermi energy) was taken to be zero.

2 Ideal Bose Gas in External Potential

Can we apply this idea to a Bose gas? Let us first consider an ideal Bose gas in an external potential, at a temperature T above the transition point. Take the potential to be harmonic:

$$V(r) = \frac{1}{2}m\omega^2 r^2 \quad (8)$$

For the Thomas-Fermi idea to be applicable, we require

$$\frac{\hbar\omega}{kT} \ll 1 \quad (9)$$

Above the transition temperature, the density is related to the fugacity $z = \exp(-\mu/kT)$ through

$$n = \frac{1}{\lambda^3} g_{3/2}(z) \quad (10)$$

where $\lambda = \sqrt{2\pi\hbar^2/mkT}$ is the thermal wavelength, and

$$g_{3/2}(z) = \sum_{\ell=1}^{\infty} \frac{z^{\ell}}{\ell^{3/2}} \quad (11)$$

This suggests that in the presence of an external potential we take the local density to be

$$n(r) = \frac{1}{\lambda^3} g_{3/2}(ze^{-\beta V(r)}) \quad (12)$$

where $\beta = 1/kT$. Integrating both sides over all space, we obtain an expression for the total number of particles:

$$N = \frac{1}{\lambda^3} \int d^3r g_{3/2}(ze^{-\beta V(r)}) \quad (13)$$

We know that $g_{3/2}(z)$ is bounded for $0 \leq z \leq 1$. So the right side is bounded. This forces Bose-Einstein condensation when N exceeds the bound. The number of atoms in the condensate N_0 is given through

$$N = N_0 + \frac{1}{\lambda^3} \int d^3r g_{3/2}(ze^{-\beta V(r)}) \quad (14)$$

In this intuitive approach, however, the Bose condensate was not described accurately.

As it turns out, the problem can be solved exactly [5]:

$$n(r) = \frac{z_1}{1 - z_1} |\psi_0(r)|^2 + \frac{1}{\lambda^3} G(z, r) \quad (15)$$

where $\psi_0(r)$ is the ground-state wave function in the potential, and

$$G(z_1, r) = \frac{1}{(2\epsilon)^{3/2}} \sum_{\ell=1}^{\infty} z_1^{\ell} \left\{ \frac{\exp[-(r/r_0)^2 \tanh(\epsilon\ell/2)]}{[1 - \exp(-2\epsilon\ell)]^{3/2}} - \exp[-(r/r_0)^2] \right\} \quad (16)$$

with

$$\begin{aligned} z_1 &= z e^{-3\hbar\omega/2kT} \\ \epsilon &= \hbar\omega/kT \\ r_0 &= \sqrt{kT/2\pi m\omega^2} \end{aligned} \tag{17}$$

The explicit occurrence of $\psi_0(r)$ shows that Bose condensation occurs in the ground state ψ_0 of the potential. The zero-momentum state is irrelevant here. The total number of particles is

$$N = \frac{z_1}{1 - z_1} + \frac{1}{\lambda^3} \int d^3r G(z, r) \tag{18}$$

which shows that the condensation is a continuous process, though it may appear to be abrupt, when N is so large that the first term can be neglected except near $z_1 = 1$.

The Thomas-Fermi approximation is good when $\epsilon \ll 1$. In that case we have

$$G(z_1, r) \approx g_{3/2} \left(z_1 e^{-\beta V(r)} \right) \tag{19}$$

and therefore

$$n(r) \approx \frac{z_1}{1 - z_1} |\psi_0(r)|^2 + \frac{1}{\lambda^3} g_{3/2} \left(z_1 e^{-\beta V(r)} \right) \tag{20}$$

which is similar to the naive formula, except for a better representation of the condensate. (The replacement of z by z_1 is inconsequential.) The lesson is that a purely intuitive approach is not satisfactory, and we need a systematic method.

3 Uniform Dilute Interacting Bose Gas

The underlying idea of the Thomas-Fermi approach is to treat a nonuniform condensate as locally uniform, with a slowly varying density. I will first review the properties of a uniform Bose gas in the dilute limit, with interparticle interactions taken into account through a scattering length $a \geq 0$.

The annihilation operator $a_{\mathbf{k}}$ of a particle of momentum $\hbar\mathbf{k}$ satisfies the commutation relation

$$[a_{\mathbf{k}}, a_{\mathbf{k}'}^\dagger] = \delta_{\mathbf{k}\mathbf{k}'} \tag{21}$$

We make a Bogolubov transformation to quasiparticle operators $\eta_{\mathbf{k}}$:

$$a_{\mathbf{k}} = x_{\mathbf{k}} \eta_{\mathbf{k}} - y_{\mathbf{k}} \eta_{-\mathbf{k}}^\dagger \quad (\mathbf{k} \neq 0) \tag{22}$$

and require that the transformation be canonical, *i.e.*

$$[\eta_{\mathbf{k}}, \eta_{\mathbf{k}'}^\dagger] = \delta_{\mathbf{k}\mathbf{k}'} \tag{23}$$

This leads to the condition

$$x_{\mathbf{k}}^2 - y_{\mathbf{k}}^2 = 1 \tag{24}$$

which can be satisfied by putting

$$\begin{aligned}x_k &= \cosh \sigma_k \\ y_k &= \sinh \sigma_k\end{aligned}\tag{25}$$

This a convenient parametrization, because interesting quantities find simple expressions:

$$\begin{aligned}\rho_k &\equiv \langle a_k^\dagger a_k \rangle = \frac{1}{2} [\cosh(2\sigma_k) - 1] \\ \Delta_k &\equiv -\langle a_k a_{-k} \rangle = \frac{1}{2} \sinh(2\sigma_k)\end{aligned}\tag{26}$$

where ρ_k measures the depletion of the unperturbed condensate:

$$N_0 = N - \sum_{\mathbf{k} \neq 0} \rho_k\tag{27}$$

and Δ_k , is a measure of off-diagonal long range order.

In the Bogolubov method, the annihilation operator for the zero-momentum state a_0 is equated to the c-number $\sqrt{N}$. Explicit solution of the problem gives

$$\tanh(2\sigma_k) = \frac{\mu}{(\hbar^2 k^2 / 2m) + \mu}\tag{28}$$

where μ is the chemical potential:

$$\mu = \frac{4\pi a \hbar^2 n}{m} \left(1 + \frac{32}{3} \sqrt{\frac{na^3}{\pi}} \right)\tag{29}$$

with n the particle density. Note that this cannot be continued to negative a ; apparently, new physics arises when the scattering length turns negative. The excitation energy of a quasiparticle of momentum $\mathbf{p}$ is given by

$$\epsilon_p = \sqrt{\left(\frac{p^2}{2m} + \mu \right)^2 - \mu^2}\tag{30}$$

4 Quasiparticle Field Operator

In the uniform case, the field operator $\Psi(\mathbf{r})$ can be put in the form

$$\Psi(\mathbf{r}) = a_0 + \psi(\mathbf{r})\tag{31}$$

with

$$\psi(\mathbf{r}) = \Omega^{-1/2} \sum_{\mathbf{k} \neq 0} a_k e^{i\mathbf{k} \cdot \mathbf{r}}\tag{32}$$

where Ω is the spatial volume. We have

$$[\psi(\mathbf{r}), \psi^\dagger(\mathbf{r}')] = \delta(\mathbf{r} - \mathbf{r}')\tag{33}$$

since a_0 is treated as a c-number. We can introduce a quasiparticle field operator:

$$\xi(\mathbf{r}) \equiv \Omega^{-1/2} \sum_{\mathbf{k} \neq 0} \eta_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{r}} \quad (34)$$

Note that the relation between ψ and ξ is non-local:

$$\psi(\mathbf{r}) = \int d^3y [X(\mathbf{x} - \mathbf{y})\xi(\mathbf{y}) - Y^*(\mathbf{x} - \mathbf{y})\xi^\dagger(\mathbf{y})] \quad (35)$$

where

$$\begin{aligned} X(\mathbf{x} - \mathbf{y}) &= \Omega^{-1/2} \sum_{\mathbf{k} \neq 0} x_{\mathbf{k}} e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} \\ Y(\mathbf{x} - \mathbf{y}) &= \Omega^{-1/2} \sum_{\mathbf{k} \neq 0} y_{\mathbf{k}} e^{i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})} \end{aligned} \quad (36)$$

For a non-uniform Bose gas, we write

$$\Psi(\mathbf{r}) = \phi(\mathbf{r}) + \psi(\mathbf{r}) \quad (37)$$

where $\phi(\mathbf{r})$ is a c-number function, such that

$$\langle \psi(\mathbf{r}) \rangle = 0 \quad (38)$$

where $\langle \rangle$ denotes ground state expectation value. We transform to quasiparticle operators by putting

$$\psi(\mathbf{r}) = \int d^3y [X(\mathbf{x}, \mathbf{y})\xi(\mathbf{y}) - Y^*(\mathbf{x}, \mathbf{y})\xi^\dagger(\mathbf{y})] \quad (39)$$

The requirement

$$[\xi(\mathbf{r}), \xi^\dagger(\mathbf{r}')] = \delta(\mathbf{r} - \mathbf{r}') \quad (40)$$

leads to the condition

$$\int d^3z [X(\mathbf{x}, \mathbf{z})X(\mathbf{z}, \mathbf{y}) - Y(\mathbf{x}, \mathbf{z})Y(\mathbf{z}, \mathbf{y})] = \delta(\mathbf{x} - \mathbf{y}) \quad (41)$$

The fulfillment of this condition in a simple fashion will guide our formulation of the Thomas-Fermi approximation.

5 Wigner Representation

In quantum mechanics, the Wigner distribution associated with a wave function $\psi(\mathbf{r})$ is defined by

$$\rho_W(\mathbf{R}, \mathbf{p}) \equiv \int d^3r \psi^*(\mathbf{R} + \mathbf{r}/2) \psi(\mathbf{R} - \mathbf{r}/2) e^{i\mathbf{p} \cdot \mathbf{r}/\hbar} \quad (42)$$

That is, we take the off-diagonal density at two different points in space, and Fourier analyze with respect to the relative separation. This is illustrated in Fig.3.




Figure 3: To get the Wigner distribution, Fourier analyze with respect to relative coordinate.

The Wigner distribution is not positive-definite, and hence not a probability; but it acts as a quasi-distribution function in phase space:

$$\begin{aligned}
(\psi, f\psi) &\equiv \int d^3r \psi^*(\mathbf{r}) f(\mathbf{r}) \psi(\mathbf{r}) = \int \frac{d^3R d^3p}{h^3} f(\mathbf{R}) \rho_W(\mathbf{R}, \mathbf{p}) \\
(\psi, \mathbf{p}\psi) &\equiv \int d^3r \psi^*(\mathbf{r}) \frac{\hbar}{i} \nabla \psi(\mathbf{r}) = \int \frac{d^3R d^3p}{h^3} \frac{\hbar}{i} \nabla \rho_W(\mathbf{R}, \mathbf{p})
\end{aligned} \tag{43}$$

For a function $X(\mathbf{x}, \mathbf{y})$ that depends on two spatial points, we define its Wigner transform as

$$X_W(\mathbf{R}, \mathbf{p}) \equiv \int d^3r X(\mathbf{R} + \mathbf{r}/2, \mathbf{R} - \mathbf{r}/2) e^{i\mathbf{p} \cdot \mathbf{r}} \tag{44}$$

with the inverse transform

$$X(\mathbf{x}, \mathbf{y}) = \int \frac{d^3p}{(2\pi)^3} e^{-i\mathbf{p} \cdot (\mathbf{x} - \mathbf{y})} X_W((\mathbf{x} + \mathbf{y})/2, \mathbf{p}) \tag{45}$$

If $C(\mathbf{x}, \mathbf{y})$ has the form

$$C(\mathbf{x}, \mathbf{y}) = \int d^3z A(\mathbf{x}, \mathbf{z}) B(\mathbf{z}, \mathbf{y}) \tag{46}$$

then its Wigner transform takes the form

$$\begin{aligned}
C_W(\mathbf{R}, \mathbf{p}) &= A_W(\mathbf{R}, \mathbf{p}) B_W(\mathbf{R}, \mathbf{p}) + \frac{1}{2i} \sum_{j=1}^3 \left(\frac{\partial A_W}{\partial R_j} \frac{\partial B_W}{\partial p_j} - \frac{\partial B_W}{\partial R_j} \frac{\partial A_W}{\partial p_j} \right) \\
&\quad - \frac{1}{8} \sum_{j=1}^3 \left(\frac{\partial^2 A_W}{\partial R_j^2} \frac{\partial^2 B_W}{\partial p_j^2} + \frac{\partial^2 B_W}{\partial R_j^2} \frac{\partial^2 A_W}{\partial p_j^2} - 2 \frac{\partial^2 A_W}{\partial R_j \partial p_j} \frac{\partial^2 B_W}{\partial R_j \partial p_j} \right) \\
&\quad + \dots
\end{aligned} \tag{47}$$

The second term is the classical Poisson bracket $\{A_W, B_W\}_{\text{PB}}$. It and the subsequent terms all depend on spatial derivatives, and would be small if the system is nearly uniform. Thus our version of Thomas-Fermi approximation consists of keeping only the first term. Errors incurred can be estimated by calculating the next non-vanishing term.

In terms of the Wigner transform, we can write

$$\int d^3z X(\mathbf{x}, \mathbf{z}) X(\mathbf{z}, \mathbf{y}) = X_W(\mathbf{R}, \mathbf{p}) X_W(\mathbf{R}, \mathbf{p}) + \frac{1}{2i} \{X_W, X_W\}_{\text{PB}} + \dots \tag{48}$$

where the second terms vanishes identically. Thus, for such an integral, errors incurred in using the Thomas-Fermi approximation starts with subsequent terms. The condition on X and Y therefore reads

$$X_W^2(\mathbf{R}, \mathbf{p}) - Y_W^2(\mathbf{R}, \mathbf{p}) \approx 1 \quad (49)$$

and is solved by setting

$$\begin{aligned} X_W(\mathbf{R}, \mathbf{p}) &= \cosh \sigma(\mathbf{R}, \mathbf{p}) \\ Y_W(\mathbf{R}, \mathbf{p}) &= \sinh \sigma(\mathbf{R}, \mathbf{p}) \end{aligned} \quad (50)$$

This make the problem very similar to the uniform case.

At zero temperature, the criterion for the validity of the Thomas-Fermi approximation is

$$\hbar\omega/\mu \ll 1 \quad (51)$$

where $\hbar\omega$ is the characteristic energy of the external potential, and μ is the chemical potential. For the dilute interacting Bose gas, μ is of order of the scattering length. Thus, the Thomas-Fermi approximation can be used only when there are interparticle interactions.

6 Variational Calculation

We study the system defined by the Hamiltonian H , with

$$H - \mu N = \int d^3x \Psi^\dagger h \Psi + \frac{1}{2} \int d^3x d^3y \Psi^\dagger \Psi^\dagger V \Psi \Psi \quad (52)$$

where $V(\mathbf{x})$ is the interparticle potential, and

$$h = -\frac{\hbar^2}{2m} \nabla^2 + V_{\text{ext}}(\mathbf{x}) - \mu \quad (53)$$

with $V_{\text{ext}}(\mathbf{x})$ the external potential. The ground state free energy is

$$F = \langle H - \mu N \rangle \quad (54)$$

where $\langle \rangle$ means expectation value with respect to the ground state of $H - \mu N$. As mentioned before, we displace the field by writing $\Psi = \phi + \psi$, where ϕ is a c-number function, such that $\langle \psi \rangle = 0$. We assume a trial form for the ground state, so that $\langle F \rangle$ has the same form as in mean-field theory, *i.e.*, we can put

$$\langle \psi^\dagger(\mathbf{y}) \psi^\dagger(\mathbf{x}) \psi(\mathbf{x}) \psi(\mathbf{y}) \rangle = \Delta^*(\mathbf{y}, \mathbf{x}) \Delta(\mathbf{x}, \mathbf{y}) + \rho(\mathbf{y}, \mathbf{x}) \rho(\mathbf{x}, \mathbf{y}) + \rho(\mathbf{y}, \mathbf{y}) \rho(\mathbf{x}, \mathbf{x}) \quad (55)$$

where

$$\begin{aligned} \rho(\mathbf{x}, \mathbf{y}) &= \langle \psi^\dagger(\mathbf{x}) \psi(\mathbf{y}) \rangle \\ \Delta(\mathbf{x}, \mathbf{y}) &= -\langle \psi(\mathbf{x}) \psi(\mathbf{y}) \rangle \end{aligned} \quad (56)$$

The ground state free energy $F[\phi, \rho, \Delta; \mu]$ is a functional of ϕ , ρ , and Δ , and also depends on μ as a parameter. The requirement $\langle \psi \rangle = 0$ means that there are no terms in F linear in ϕ .

Although we do not need the trial state explicitly, it can be explicitly constructed if desired. One can show that the wave functional of this state is of Gaussian form [8]. Thus, we have a true variational problem.

We rewrite the functions ρ and Δ in $F[\phi, \rho, \Delta; \mu]$ in terms of their Wigner transforms, and implement our version of the Thomas-Fermi approximation, as explained before. We transform to quasiparticle field operators, and find that, as in the uniform case, ρ_W and Δ are parametrized by a single function:

$$\begin{aligned}\rho_W(\mathbf{R}, \mathbf{p}) &= \frac{1}{2}[\cosh(2\sigma(\mathbf{R}, \mathbf{p})) - 1] \\ \Delta_W(\mathbf{R}, \mathbf{p}) &= \frac{1}{2} \sinh(2\sigma(\mathbf{R}, \mathbf{p}))\end{aligned}\tag{57}$$

The Free energy reduces to the form $F = \int d^3R f(\mathbf{R})$. We obtain equations for σ and ϕ by minimizing F . The equation for ϕ is a modified Gross-Pitaevskii or non-linear Schrödinger equation (NLSE):

$$\left[-\frac{\hbar^2}{2m} \nabla^2 + V_{\text{ext}}(\mathbf{r}) + U(\mathbf{r}) - \mu + v(0)\phi^2(\mathbf{r}) \right] \phi(\mathbf{r}) = 0\tag{58}$$

where $U(\mathbf{r})$ is a self-consistent potential that depends on σ . It is unimportant for low densities.

7 Dilute Interacting Gas in Harmonic Trap

I will just quote some results for a dilute gas in a harmonic trap. The external potential is

$$V_{\text{ext}} = \frac{\hbar\omega}{2} \left(\frac{r}{L} \right)^2\tag{59}$$

For particles of mass m ,

$$L = \sqrt{\hbar/m\omega}\tag{60}$$

which is the extend of the ground state wave function. For the interparticle interaction, we use a pseudopotential

$$\frac{4\pi a \hbar^2}{m} \delta(\mathbf{r}) \frac{\partial}{\partial r} r\tag{61}$$

The sole effect of the differential operator above is the removal of a divergence in the ground state energy. The three important lengths in the problem are

$$\begin{array}{ll} L & \text{(Extend of ground state wave function)} \\ a & \text{(Scattering length)} \\ R_0 & \text{(Extend of condensate)} \end{array}\tag{62}$$

They are illustrated in Fig.4.




Figure 4: Length scales in atomic trap. Groundstate wave function is $\psi_0(r)$. Condensate wave function is $\phi_0(r)$.

For low densities, the non-linear coupled equations for σ and ϕ are solved by iteration, and one iteration suffices. The chemical potential is found to be

$$\mu = \frac{\hbar\omega}{2} \left(\frac{15a}{2L} \right)^{2/5} \left[1 + \frac{\sqrt{2}}{60} \left(\frac{15a}{L} \right)^{6/5} N^{1/5} \right] \quad (63)$$

The requirement $\hbar\omega/\mu \ll 1$ means

$$\frac{L}{aN} \ll 1 \quad (64)$$

The extend of the condensate is given by

$$\frac{R_0}{L} = \left(\frac{15aN}{L} \right)^{1/5} \quad (65)$$

For the method to be valid we must have $R_0 \gg L$. By neglecting the term $\nabla^2\phi$ in the NLSE, we find

$$\phi^2(r) = \frac{R_0^2}{8\pi a L^4} \left[1 - \left(\frac{r}{R_0} \right)^2 \right] \quad (66)$$

In Fig.5 we show the shape of the condensate and estimated errors. Fig.5(a) shows $\phi(r)$ as a function of r in units of L , for $N=10^3$, and 10^6 .

Fig.5(b) shows the errors arising from the neglect of $\nabla^2\phi$. This is “trivial,” as it can be corrected through numerical computation.

Fig.5(c) shows the errors incurred due to the Thomas-Fermi approximation, and are intrinsic to the method. They are small except at the edge of the condensate.




Figure 5: (a) Condensate wave functions for $N = 10^3$ and 10^6 ; (b) Error incurred in neglecting kinetic term in NLSE; (c) Error incurred in Thomas-Fermi approximation. Length scale on horizontal axis is in units of L , the extend of the groundstate wave function. Calculations are done for $L = 10^{-4} \text{ cm}$, scattering length $= 5 \times 10^{-7} \text{ cm}$.

8 Quasiparticle Excitation

The local excitation energy should be measured from the chemical potential:

$$\epsilon_p(r) = \mu + \sqrt{\left[\frac{p^2}{2m} + \mu_{\text{eff}}(r)\right]^2 - \mu_{\text{eff}}^2(r)} \quad (67)$$

where

$$\mu_{\text{eff}}(r) = \mu - V_{\text{ext}}(r) \quad (68)$$

It describes a phonon with a position-dependent sound velocity. The excitation energy density of the system is given by

$$g(\epsilon) = \sum_i \delta(\epsilon - \epsilon_i) \quad (69)$$

where ϵ_i is the energy of the i th excited state. In the spirit of the Thomas-Fermi approximation, we take

$$g(\epsilon) = \int \frac{d^3r d^3p}{h^3} \delta(\epsilon - \epsilon_p(r)) \quad (70)$$

The results for $N = 10^3$ and 10^6 are shown in Fig.6 and Fig.7, with comparison to the ideal gas.

Further details can be found in [7].

This work was supported in part by funds provided by the U.S. Department of Energy under cooperative agreement # DE-FC02-94ER40818.

Density of States for 10^3 particles

$L = 10^4$ cm $a = 5 \times 10^{-7}$ cm




Figure 6: Density of states in harmonic trap, for $N = 10^3$.

Density of States for 10^6 particles

$L = 10^4 \text{ cm}$ $a = 5 \times 10^{-7} \text{ cm}$




Figure 7: Same as Fig.6, but for $N = 10^6$.

References

- [1] L.H. Thomas, Proc. Camb. Phil. Soc. **23** 542 (1927).
- [2] E. Fermi, Atti Rentes Accad. Naz. Lincei Rend. Cl. Sci. Fis, Mat. Natur.,**6**. 602 (1927).
- [3] E. Lieb and B. Simon, Phys. Rev. Lett. **31**, 681 (1973).
- [4] V.V. Goldman, I.F. Silvera, and A.J. Legget, Phys. Rev. **B24**, 2870 (1981).
- [5] J. Oliva, Phys. Rev, **B39**, 4197 (1989).
- [6] T.T Chou, C.N. Yang, and L.W. Yu, Stonybrook preprints (1996).
- [7] E. Timmermans, P. Tommasini, and K. Huang, (to be published).
- [8] P. Tommasini and K. Huang, J. Res. NIST (August, 1996).